

August 2019

Dear Valued Customer,

The first registered visibly distinct skin antiseptic range'

We are pleased to advise that the Surgi-Prep® range will soon be available, bringing multiple benefits for your patients, clinicians and organisation, including:

- **Better identification**, with visually distinct solutions non-skin-staining pink or skin-staining red'
- **Broad spectrum efficacy** including multi resistant *S.aureus* and multi resistant *E.colli*²
- **Choice of volume and application options** to suit different healthcare environments'
- **Pourable format** to meet individual patient needs, **with less waste**

We will notify you as soon as stock is available and provide full details of pricing. Please see overleaf for product information and product codes.

Meanwhile, if you have any questions about our products or the launch, our Customer Service Team will be pleased to assist you. They can be contacted on:

T **1800 805 546** or E customerservice@perrigo.com.au

Yours faithfully



Stephan Bousias
Associate Director - Hospital & Specialty



VISIBLE RISK REDUCTION FOR A BROAD RANGE OF SKIN ANTISEPSIS NEEDS^{1,3}

The Surgi-Prep® range offers unprecedented flexibility and choice to help clinicians 'do what's right for the site'.

Surgi-Prep® C Chlorhexidine 0.5% and Ethanol 70%

Surgi-Prep® C+ Chlorhexidine 2% and Ethanol 70%

PRODUCT DESCRIPTION **PRODUCT CODE** **ARTG No.**

Surgi-Prep® C Red 30mL 5000023484 319263

Surgi-Prep® C Red 100mL 5000023485 319263

Surgi-Prep® C Pink 30mL 5000023478 319262

Surgi-Prep® C Pink 100mL 5000023479 319262

Surgi-Prep® C Pink 500mL 5000023480 319262

PRODUCT DESCRIPTION **PRODUCT CODE** **ARTG No.**

Surgi-Prep® C+ Red 30mL 5000023487 319265

Surgi-Prep® C+ Red 100mL 5000023488 319265

Surgi-Prep® C+ Pink 30mL 5000023481 319264

Surgi-Prep® C+ Pink 100mL 5000023482 319264

Surgi-Prep® C+ Pink 500mL 5000023483 319264

ALWAYS READ THE LABEL. FOLLOW DIRECTIONS FOR USE.

References: 1. Australia Register of Therapeutic Goods AUST R 319262, 319263, 319264 & 319265. 2019. 2. Perrigo Data on file. 3. Australia and New Zealand College of Anaesthetists (ANZCA). Topical application of chlorhexidine and the risks of accidental injection in regional anaesthesia and vascular access procedures. 2018. Available at: <http://www.anzca.edu.au/documents/8-2-2-anzca-and-acsqhc-joint-statement-on-chlorhex.pdf> Last accessed: 1 August 2019.

Surgi-Prep® is a registered trademark of Perrigo Australia. ABN 56 009 293 136. 25–29 Delawney Street, Balcatta, Western Australia, 6021, AUSTRALIA. <https://www.perrigo.com.au>. For Medical Information enquiries: 1800 805 546 or Medicalinfo@perrigo.com.au. To report an adverse event: 1800 805 546, <https://www.perrigo.com.au/contact/>. 201908-01861V1. August 2019. 15715.



Surgi-Prep® C Red Chlorhexidine 0.5% and Ethanol 70%

100mL

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.

WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissue. Avoid getting solution into eyes. Wet hair is flammable. Flammable. Flammable. Flammable. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs, avoid contact with eyes. Contains ethanol 70% v/v.

DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.

OTHER INFORMATION: Protect from light. Store below 25°C.

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

Perrigo
Surgi-Prep® C red
Chlorhexidine gluconate 0.5% w/v
and Ethanol 70% v/v

100mL Application AUST R 319263
Skin Staining

Perrigo Australia, 25-29 Delawney Street, Perrigo
Balaclava, VIC 3207 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

DANGER Highly flammable liquid and vapour. Causes serious eye irritation.

9 431991 2 035545

EXP 5000023485 1000011780



30mL

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.

WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissue. Solution is flammable. Flammable. Flammable. Flammable. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs, avoid contact with eyes. Contains ethanol 70% v/v.

DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.

OTHER INFORMATION: Protect from light. Store below 25°C.

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

Perrigo
Surgi-Prep® C red
Chlorhexidine gluconate 0.5% w/v
and Ethanol 70% v/v

30mL Application AUST R 319263
Skin Staining

Perrigo Australia, 25-29 Delawney Street, Perrigo
Balaclava, VIC 3207 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

DANGER Highly flammable liquid and vapour.

9 431991 2 035530

EXP 5000023484 1000011779



Surgi-Prep® C+ Red Chlorhexidine 2% and Ethanol 70%



INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.

WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissue. Wet hair is flammable. Flammable. Flammable. Flammable. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs, avoid contact with eyes. Contains ethanol 70% v/v.

DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.

OTHER INFORMATION: Protect from light. Store below 25°C.

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

Perrigo
Surgi-Prep® C+ red
Chlorhexidine gluconate 2% w/v
and Ethanol 70% v/v

100mL Application AUST R 319265
Skin Staining

Perrigo Australia, 25-29 Delawney Street, Perrigo
Balaclava, VIC 3207 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

DANGER Highly flammable liquid and vapour. Causes serious eye irritation.

9 431991 2 035576

EXP 5000023488 1000011783

100mL



INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.

WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissue. Solution is flammable. Flammable. Flammable. Flammable. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs, avoid contact with eyes. Contains ethanol 70% v/v.

DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.

OTHER INFORMATION: Protect from light. Store below 25°C.

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

Perrigo
Surgi-Prep® C+ red
Chlorhexidine gluconate 2% w/v
and Ethanol 70% v/v

30mL Application AUST R 319265
Skin Staining

Perrigo Australia, 25-29 Delawney Street, Perrigo
Balaclava, VIC 3207 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

DANGER Highly flammable liquid and vapour.

9 431991 2 035569

EXP 5000023487 1000011782

30mL

Surgi-Prep® C Pink

Chlorhexidine 0.5% and Ethanol 70%

Surgi-Prep® C+ Pink

Chlorhexidine 2% and Ethanol 70%

500mL

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

DANGER
Highly flammable liquid and vapour. Causes serious eye irritation.
ETHANOL SOLUTION, UN1170 Class 3 PGII • 2YE

Perrigo
Surgi-Prep® C pink
Chlorhexidine gluconate 0.5% w/v and Ethanol 70% v/v

500mL Application AUST R 319262

Perrigo Australia, 25-29 Delawney Street, Balakata, WA 6021 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

9 319912 03 5491

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.
WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissues. Avoid getting solution into hairy areas. Wet hair is flammable. Hair may take up to 1 hour to dry. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs. Avoid contact with eyes. Contains ethanol 70% v/v.
DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.
OTHER INFORMATION: Protect from light. Store below 25°C.

EXP 5000023480 1000011775



CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

DANGER
Highly flammable liquid and vapour. Causes serious eye irritation.
ETHANOL SOLUTION, UN1170 Class 3 PGII • 2YE

Perrigo
Surgi-Prep® C+ pink
Chlorhexidine gluconate 2% w/v and Ethanol 70% v/v

500mL Application AUST R 319264

Perrigo Australia, 25-29 Delawney Street, Balakata, WA 6021 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

9 319912 03 5521

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.
WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissues. Avoid getting solution into hairy areas. Wet hair is flammable. Hair may take up to 1 hour to dry. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs. Avoid contact with eyes. Contains ethanol 70% v/v.
DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.
OTHER INFORMATION: Protect from light. Store below 25°C.

EXP 5000023483 1000011778

500mL

100mL

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

DANGER
Highly flammable liquid and vapour. Causes serious eye irritation.

Perrigo
Surgi-Prep® C pink
Chlorhexidine gluconate 0.5% w/v and Ethanol 70% v/v

100mL Application AUST R 319262

Perrigo Australia, 25-29 Delawney Street, Balakata, WA 6021 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

9 319912 10 35484

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.
WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissues. Avoid getting solution into hairy areas. Wet hair is flammable. Hair may take up to 1 hour to dry. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs. Avoid contact with eyes. Contains ethanol 70% v/v.
DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.
OTHER INFORMATION: Protect from light. Store below 25°C.

EXP 5000023479 1000011774



CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

DANGER
Highly flammable liquid and vapour. Causes serious eye irritation.

Perrigo
Surgi-Prep® C+ pink
Chlorhexidine gluconate 2% w/v and Ethanol 70% v/v

100mL Application AUST R 319264

Perrigo Australia, 25-29 Delawney Street, Balakata, WA 6021 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

9 319912 10 35514

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.
WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissues. Avoid getting solution into hairy areas. Wet hair is flammable. Hair may take up to 1 hour to dry. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs. Avoid contact with eyes. Contains ethanol 70% v/v.
DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.
OTHER INFORMATION: Protect from light. Store below 25°C.

EXP 5000023482 1000011777

100mL

30mL

CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

DANGER
Highly flammable liquid and vapour.

Perrigo
Surgi-Prep® C pink
Chlorhexidine gluconate 0.5% w/v and Ethanol 70% v/v

30mL Application AUST R 319262

Perrigo Australia, 25-29 Delawney Street, Balakata, WA 6021 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

9 319912 10 35477

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.
WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissues. Solution is flammable, allow to dry completely before proceeding. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs. Avoid contact with eyes. Contains ethanol 70% v/v.
DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.
OTHER INFORMATION: Protect from light. Store below 25°C.

EXP 5000023478 1000011773



CAUTION
KEEP OUT OF REACH OF CHILDREN
DO NOT SWALLOW

DANGER
Highly flammable liquid and vapour.

Perrigo
Surgi-Prep® C+ pink
Chlorhexidine gluconate 2% w/v and Ethanol 70% v/v

30mL Application AUST R 319264

Perrigo Australia, 25-29 Delawney Street, Balakata, WA 6021 Australia. Phone: 1800 805 546
Distributed in New Zealand by Orion Laboratories (NZ) Ltd

9 319912 10 35507

INDICATIONS: Skin antiseptic applied prior to invasive medical procedures.
WARNINGS: Avoid direct contact with neural tissues such as meninges, spinal cord and brain tissues. Solution is flammable, allow to dry completely before proceeding. Do not use on open skin wounds, broken or damaged skin. This product contains chlorhexidine. Severe allergic reactions can occur. Stop use if this occurs. Avoid contact with eyes. Contains ethanol 70% v/v.
DIRECTIONS FOR USE: Clean the application site if visibly soiled. Apply solution to intact skin for at least 30 seconds on dry sites and 2 minutes on moist sites, using repeated circular strokes, working out from the centre. Do not blot or wipe away. For detailed instructions see www.perrigo.com.au.
OTHER INFORMATION: Protect from light. Store below 25°C.

EXP 5000023481 1000011776

30mL

Surgi-Prep[®]



ALWAYS READ THE LABEL. FOLLOW DIRECTIONS FOR USE.

Surgi-Prep[®] is a registered trademark of Perrigo Australia. ABN 56 009 293 136. 25–29 Delawney Street, Balcatta, Western Australia, 6021, AUSTRALIA. <https://www.perrigo.com.au>. For Medical Information enquiries: 1800 805 546 or Medicalinfo@perrigo.com.au. To report an adverse event: 1800 805 546, <https://www.perrigo.com.au/contact/>. 201908-01861V1. August 2019. 15715.

